

# A GENERALIZED POINCARÉ-LELONG FORMULA

MATS ANDERSSON

**ABSTRACT.** We prove a generalization of the classical Poincaré-Lelong formula. Given a holomorphic section  $f$ , with zero set  $Z$ , of a Hermitian vector bundle  $E \rightarrow X$ , let  $S$  be the line bundle over  $X \setminus Z$  spanned by  $f$  and let  $Q = E/S$ . Then the Chern form  $c(D_Q)$  is locally integrable and closed in  $X$  and there is a current  $W$  such that  $dd^c W = c(D_E) - c(D_Q) - M$ , where  $M$  is a current with support on  $Z$ . In particular, the top Bott-Chern class is represented by a current with support on  $Z$ . We discuss positivity of these currents, and we also reveal a close relation with principal value and residue currents of Cauchy-Fantappiè-Leray type.

## 1. INTRODUCTION

Let  $f$  be a holomorphic (or meromorphic) section of a Hermitian line bundle  $L \rightarrow X$ , and let  $[Z]$  be the current of integration over the divisor  $Z$  defined by  $f$ . The Poincaré-Lelong formula states that

$$dd^c \log(1/|f|) = c_1(D_L) - [Z],$$

where  $c_1(D_L)$  is the first Chern form associated with the Chern connection  $D_L$  on  $L$ , i.e.,  $c_1(D_L) = \kappa \Theta_L$ , where  $\Theta_L$  is the curvature; here and throughout this paper  $\kappa = i/2\pi$  and  $d^c = \kappa(\bar{\partial} - \partial)$  so that

$$dd^c = \frac{i}{\pi} \partial \bar{\partial} = 2\kappa \partial \bar{\partial}.$$

If  $U$  is the meromorphic section of the dual bundle  $L^*$  such that  $U \cdot f = 1$ , then  $R = \bar{\partial}U$  is a  $(0, 1)$ -current, and we have the global factorization

$$(1.1) \quad [Z] = R \cdot D_L f / 2\pi i.$$

If  $A = -2\kappa \partial \log(1/|f|)$ , then clearly  $dA = \bar{\partial}A = c_1(D_L) - [Z]$ , and it is easily checked that  $A = U \cdot D_L f / 2\pi i$ . In this paper we consider analogous formulas for a holomorphic section  $f$  of a higher rank bundle, and our main result is the following generalization of the Poincaré-Lelong formula.

**Theorem 1.1.** *Let  $f$  be a holomorphic section of the Hermitian vector bundle  $E \rightarrow X$  of rank  $m$ . Let  $Z = \{f = 0\}$ , let  $S$  denote the (trivial) line bundle over  $X \setminus Z$  generated by  $f$ , and let  $Q = E/S$ , equipped with the induced Hermitian metric.*

---

*Date:* March 16, 2010.

The author was partially supported by the Swedish Natural Science Research Council.

(i) The Chern form  $c(D_Q)$  is locally integrable in  $X$  and its natural extension to  $X$  is closed. Moreover, the forms  $\log |f|c(D_Q)$  and

$$(1.2) \quad |f|^{2\lambda} \frac{\Re \partial |f|^2 \wedge \bar{\partial} |f|^2}{|f|^4} \wedge c(D_Q), \quad \lambda > 0,$$

are locally integrable in  $X$ , and

$$(1.3) \quad M = \lim_{\lambda \rightarrow 0^+} \lambda |f|^{2\lambda} \frac{\Re \partial |f|^2 \wedge \bar{\partial} |f|^2}{|f|^4} \wedge c(D_Q) = dd^c(\log |f|c(D_Q)) \mathbf{1}_Z$$

is a closed current of order zero with support on  $Z$ . If  $\text{codim } Z = p$ , then

$$M = M_p + M_{p+1} \cdots + M_{\min(m,n)},$$

where  $M_k$  has bidegree  $(k, k)$ , and

$$M_p = \sum \alpha_j [Z_j^p],$$

where  $Z_j^p$  are the irreducible components of codimension precisely  $p$ , and  $\alpha_j$  are the Hilbert-Samuel multiplicities of  $f$ .

(ii) There is a current  $W$  of bidegree  $(*, *)$  and order zero in  $X$  which is smooth in  $X \setminus Z$ , and with logarithmic singularity at  $Z$ , such that

$$(1.4) \quad dd^c W = c(D_E) - C(D_Q) - M,$$

where  $C(D_Q)$  denote the natural extension of  $c(D_Q)$ .

Here  $c(D)$  denotes the Chern form with respect to the Chern connection  $D$  associated to the Hermitian structure, i.e.,  $c(D) = \det(\Re \Theta + I)$ , where  $\Theta = D^2$  is the curvature tensor. We let  $c_k(D)$  denote the component of bidegree  $(k, k)$ .

For an explicit expression for  $W$ , see Definition 1 in Section 4. If  $W_k$  denotes the component of bidegree  $(k, k)$ , then (1.4) means that

$$(1.5) \quad dd^c W_{k-1} = c_k(D_E) - c_k(D_Q) - M_k.$$

Since  $Q$  has rank  $m - 1$ ,  $c_m(D_Q) = 0$ , and therefore

$$dd^c W_{m-1} = c_m(D_E) - M_m,$$

which means that the current  $M_m$  represents the top degree Bott-Chern class  $\hat{c}_m(E)$ . It also follows that the Bott-Chern class  $\hat{c}_k(E)$  is equal to  $\hat{c}_k(Q)$  if  $k < p$ .

If  $E$  is a line bundle, then, see Definition 1,  $W = W_0 = \log(1/|f|)$ , so (1.5) is then the usual Poincaré-Lelong formula.

In [8] Bott and Chern developed a method of transgression which in particular gives a form  $w$  in  $X \setminus Z$  such that  $dd^c w = c(D_E) - c(D_Q)$ . It is not unexpected that one can extend this construction across  $Z$  by a careful analysis of the occurring singularities at  $Z$ . In the recent paper [17], Meo proves (1.5) for  $k = p$ . Previously this formula was proved in [7] in the case when  $f$  defines a complete intersection, i.e.,  $p = m$ . A

variety of analogous formulas for  $d$  rather than  $dd^c$  are constructed in quite general (non-holomorphic) situations in [12], [13], [14], and [15].

Clearly  $M_p$  is always a positive current. It follows from (1.3) that  $M_k$  is positive if  $c_{k-1}(D_Q)$  is a positive form. For an even more precise formula for  $M$ , see Proposition 7.5.

Let us say that  $E$  is positive if  $E^*$  is Nakano negative.

**Theorem 1.2.** *Assume that  $E$  is positive. Then  $c(D_E)$  is a positive form,  $C(D_Q)$  and  $M$  are positive currents, and (one can choose  $W$  such that)  $W$  is positive where  $|f| \leq 1$ .*

If  $A = -2\kappa\partial W$  we have, cf., (1.4),

$$(1.6) \quad \bar{\partial}A = dA = c(D_E) - c(D_Q) - M.$$

In [1] we introduced a residue current  $R = R_p + \cdots + R_{\min(m,n)}$ , associated with  $f$ , with support on  $Z$ , where  $R_k$  is a  $(0, k)$ -current with values in  $\Lambda^k E^*$ , and a principal value current  $U = U_1 + \cdots + U_m$  such that  $(\delta_f - \bar{\partial})U = 1 - R$ , where  $\delta_f$  denotes contraction with  $f$ . When  $E$  is a line bundle, then  $U = 1/f$  and  $R = \bar{\partial}(1/f)$ . In analogy to (1.1) we can factorize  $M_p$  as

$$M_p = R_p \cdot (D_E f)^p / p!;$$

this was proved in [2]. We have a similar, but somewhat more involved, formula for the whole current  $M$ , see (6.4) in Section 6. In a similar way we can express  $A$  and  $c(D_Q)$ , see (6.5) and (6.6), in terms of the current  $U$ .

*Remark 1.* Let  $f_1, \dots, f_r$  be holomorphic sections of  $E$  and let  $Z$  be the analytic set where they are linearly dependent. Moreover, let  $S$  be the trivial rank  $r$ -subbundle of  $E$  over  $X \setminus Z$  generated by  $f_j$  and let  $Q = E/S$ . Then  $c(D_Q)$  has a natural current extension  $C(D_Q)$  across  $Z$  and there is a closed current  $M$  of bidegree  $(*, *)$  with support on  $Z$  and a current  $A$  such that

$$(1.7) \quad dA = \bar{\partial}A = c(D_E) - C(D_Q) - M.$$

This can be proved by a small modification of the argument in this paper; in the case  $Z$  has generic dimension such a formula was proved already in [14], and the general case should be contained in [15]. It follows from (1.7) that the current  $M_k$  is a representative for  $c_k(D_E)$  for  $k > m - r$ .

However, we have no analogous formula for  $dd^c$ . □

As indicated above, the proof of Theorem 1.1 relies on the construction in [8], combined with a careful control of the singularities at  $Z$ . To begin with one constructs a form  $v$  in  $X \setminus Z$  such that

$$dd^c v = c(D_E) - c(D_S)c(D_Q).$$

By Hironaka's theorem and toric resolutions, following [4] and [18], we can prove that this equality has meaning in the current sense across  $Z$ . Here a crucial point is an explicit formula for the Chern form  $c(D_Q)$  (Proposition 4.2) from which it is easy to conclude that  $c(D_Q)$  has a smooth extension across the singularity after an appropriate blow-up. By the usual Poincaré-Lelong formula,  $c(D_S) - 1 = dd^c \log(1/|f|)$  outside  $Z$ , and we can conclude that (1.4) holds (if the capitals denote the natural extensions across  $Z$ ) with

$$W = \log(1/|f|)C(D_Q) - V,$$

and  $M = dd^c(\log |f|C(D_Q))\mathbf{1}_Z$ . Theorem 1.2 follows essentially by applying ideas in [8].

In Section 7 we discuss the positivity and prove Theorem 1.2, essentially by applying ideas from [8]. The paper is concluded by some examples.

## 2. PRELIMINARIES

We first recall the differential geometric definition of Chern classes. Let  $E \rightarrow X$  be any differentiable complex vector bundle over a differential manifold  $X$ , with connection  $D: \mathcal{E}_k(X, E) \rightarrow \mathcal{E}_{k+1}(X, E)$  and curvature tensor  $D^2 = \Theta \in \mathcal{E}_2(X, \text{End}E)$ . The connection  $D = D_E$  induces in a natural way a connection  $D_{\text{End}E}$  on the bundle  $\text{End}E$  by the formula  $Dg \cdot \xi = D(g \cdot \xi) - g \cdot D\xi$ , and in a similar way there is a natural connection  $D_{E^*}$  on the dual bundle  $E^*$ , etc. In particular we have Bianchi's identity

$$(2.1) \quad D_{\text{End}E}\Theta = 0.$$

If  $I$  denotes the identity mapping on  $E$ , then  $c(D) = \det(\kern-0.1em\mathfrak{s}\Theta + I)$  is a welldefined differential form whose terms have even degrees, which is called the Chern form of  $D$ . It is a basic fact that  $c(D)$  is a closed form. Moreover its de Rham cohomology class is independent of  $D$  and is called the (total) Chern class  $c(F)$  of the bundle  $F$ .

To prove this, one can consider a smooth one-parameter family  $D_t$  of connections of  $F$  with  $D_0 = D$ . If  $E'$  is the pull-back of  $E$  to  $X \times [0, 1]$ , then  $D' = D_t + d_t$  is a connection on  $E'$  and its curvature tensor is

$$\Theta' = \Theta_t + dt \wedge \dot{D}_t$$

where  $\dot{D}_t = dD_t/dt$ . It is readily checked that it is an element in  $\mathcal{E}_1(X, \text{End}(F))$ . Since  $(d + d_t) \det(\kern-0.1em\mathfrak{s}\Theta' + I) = 0$  we have that

$$d_\zeta \int_0^1 \det(\kern-0.1em\mathfrak{s}\Theta' + I) = - \int_0^1 d_t \det(\kern-0.1em\mathfrak{s}\Theta' + I) = c(D) - c(D_1).$$

In order to make the computation more explicit we introduce the exterior algebra bundle  $\Lambda = \Lambda(T^*(X) \oplus F \oplus F^*)$ . Any section  $\xi \in \mathcal{E}_k(X, F)$  corresponds to a section  $\tilde{\xi}$  of  $\Lambda$ ; if  $\xi = \xi_1 \otimes e_1 + \dots + \xi_m \otimes e_m$

in a local frame  $e_j$ , then we let  $\tilde{\xi} = \xi_1 \wedge e_1 + \dots + \xi_m \wedge e_m$ . In the same way,  $a \in \mathcal{E}_k(X, \text{End} E)$  can be identified with

$$\tilde{a} = \sum_{jk} a_{jk} \wedge e_j \wedge e_k^*,$$

if  $e_j^*$  is the dual frame, and  $a = \sum_{jk} a_{jk} \otimes e_j \otimes e_k^*$  with respect to these frames. A given connection  $D = D_F$  on  $F$  extends in a unique way to a linear mapping  $\mathcal{E}(X, \Lambda) \rightarrow \mathcal{E}(X, \Lambda)$  which is an anti-derivation with respect to the wedge product in  $\Lambda$ , and such that it acts as the exterior differential  $d$  on the  $T^*(X)$ -factor. It is readily seen that

$$\widetilde{D_E \xi} = D \tilde{\xi},$$

if  $\xi$  is a form-valued section of  $E$ . In the same way we have

**Lemma 2.1.** *If  $a \in \mathcal{E}_k(X, \text{End} E)$ , then*

$$(2.2) \quad \widetilde{D_{\text{End} E} a} = D \tilde{a}.$$

*Proof.* If  $\xi \in \mathcal{E}_k(X, E)$  and  $\eta \in \mathcal{E}(X, E^*)$ , then

$$D_{\text{End} E}(\xi \otimes \eta) = D_E \xi \otimes \eta + (-1)^k \xi \otimes D_{E^*} \eta,$$

and thus the snake of  $D_{\text{End} E}(\xi \otimes \eta)$  is equal to

$$\widetilde{D_E \xi} \wedge \eta + (-1)^{k+1} \tilde{\xi} \wedge \widetilde{D_{E^*} \eta} = D(\tilde{\xi} \wedge \eta)$$

as claimed.  $\square$

Since  $D_{\text{End} E} I = 0$ , ( $I = I_E$ ) we have from (2.1) and Lemma 2.1 that

$$(2.3) \quad D \tilde{\Theta} = 0 \quad \text{and} \quad D \tilde{I} = 0.$$

We let  $\tilde{I}_m = \tilde{I}^m / m!$  and use the same notation for other forms in the sequel. Any form  $\omega$  with values in  $\Lambda$  can be written  $\omega = \omega' \wedge \tilde{I}_m + \omega''$  uniquely, where  $\omega''$  has lower degree in  $e_j, e_k^*$ . If we define

$$\int_e \omega = \omega',$$

then this integral is of course linear and moreover

$$(2.4) \quad d \int_e \omega = \int_e D \omega.$$

In fact, since  $D \tilde{I}_m = 0$ ,

$$\int_e D \omega = \int_e d \omega' \wedge \tilde{I}_m + D \omega'' = d \omega' = d \int_e \omega.$$

Observe that

$$(2.5) \quad c(D) = \int_e (\lrcorner \tilde{\Theta} + \tilde{I})_m = \int_e e^{\lrcorner \tilde{\Theta} + \tilde{I}}.$$

Lemma 2.1 and (2.3) together imply that the Chern form  $c(D)$  is closed. Furthermore, following the outline above, we get the formula

$$(2.6) \quad d \int_0^1 \int_e \tilde{\mathbb{N}}\tilde{D} \wedge e^{\mathbb{N}\tilde{\Theta}_t + \tilde{I}} = c(D_1) - c(D_0),$$

thus showing that  $c(D_0)$  and  $c(D_1)$  are cohomologous.

Recall that if the connection  $D$  is modified to  $D_1 = D - \gamma$ , where  $\gamma \in \mathcal{E}_1(X, \text{End}E)$ , then  $\Theta_1 = \Theta - D_{\text{End}E}\gamma + \gamma \wedge \gamma$ . If we form the explicit homotopy  $D_t = D - t\gamma$ , therefore

$$(2.7) \quad \Theta_t = \Theta - tD_{\text{End}E}\gamma + t^2\gamma \wedge \gamma$$

and hence, by Lemma 2.1,

$$(2.8) \quad \tilde{\Theta}_t = \tilde{\Theta} - tD\tilde{\gamma} + t^2\widetilde{\gamma \wedge \gamma}.$$

### 3. BOTT-CHERN CLASSES

From now on we assume that  $E$  is a holomorphic Hermitian bundle and that  $D_E$  is the Chern connection and  $D'_E$  is its  $(1,0)$ -part. Then the induced connection  $D_{E^*}$  on  $E^*$  is the Chern connection on  $E^*$  etc. In particular, our mapping  $D$  on  $\Lambda$  is of type  $(1,0)$ , i.e.,  $D = D' + \bar{\partial}$ .

Let  $E \rightarrow X$  be a Hermitian vector bundle with Chern connection  $D_E$ . The Bott-Chern class  $\hat{c}(E)$  is the equivalence class of the Chern form  $c(D_E)$  in

$$\frac{\oplus_k \mathcal{E}_{k,k}(X) \cap \text{Ker } d}{\oplus_k dd^c \mathcal{E}_{k,k}(X)}.$$

**Lemma 3.1.** *Let  $D$  be a connection depending smoothly on a real parameter  $t$ . Moreover, assume that  $L \in \mathcal{E}(X, \text{End}(E))$  depends smoothly on  $t$  and that*

$$(3.1) \quad D'_{\text{End}E}L = \dot{D}.$$

*Also assume that  $\Theta_t$  has bidegree  $(1,1)$  for all  $t$ . If*

$$v = -\frac{1}{2} \int_0^1 \int_e \tilde{L}_t \wedge e^{\mathbb{N}\tilde{\Theta}_t + \tilde{I}} dt,$$

*then  $-2\mathbb{N}\partial v = b$ , where*

$$b = \int_0^1 \int_e \tilde{\mathbb{N}}\tilde{D}_t \wedge e^{\mathbb{N}\tilde{\Theta}_t + \tilde{I}} dt.$$

This lemma as well as the other material in this section is taken from [8]. However, we use a somewhat different formalism, and for the reader's convenience we supply some simple proofs.

*Proof.* In view of (2.4) we have that (suppressing the index  $t$ )

$$d \int_e \tilde{L} \wedge e^{\mathbb{N}\tilde{\Theta} + \tilde{I}} = \int_e D\tilde{L} \wedge e^{\mathbb{N}\tilde{\Theta} + \tilde{I}},$$

and by identifying bidegrees we get that

$$\partial \int_e \tilde{L} \wedge e^{\aleph \tilde{\Theta} + \tilde{I}} = \int_e D' \tilde{L} \wedge e^{\aleph \tilde{\Theta} + \tilde{I}} = \int_e \tilde{D} \wedge e^{\aleph \tilde{\Theta} + \tilde{I}}.$$

□

Since  $db = c(D_1) - c(D_0)$ , cf., (2.6), we thus have

$$(3.2) \quad -dd^c v = c(D_1) - c(D_0).$$

By deforming the metric one can use this lemma to show that  $\hat{c}(E)$  is independent of the Hermitian structure on  $E$ , see [8]. However we are interested in a somewhat different situation. Assume that we have the short exact sequence of Hermitian vector bundles

$$(3.3) \quad 0 \rightarrow S \xrightarrow{j} E \xrightarrow{g} Q \rightarrow 0,$$

where  $Q$  and  $S$  are equipped with the metrics induced by the Hermitian metric of  $E$ . Then

$$(3.4) \quad j^* \oplus g: E \rightarrow S \oplus Q$$

is a smooth vector bundle isomorphism. If  $D_S$  and  $D_Q$  are the Chern connections on  $S$  and  $Q$  respectively, then

$$(3.5) \quad D_E \sim \begin{pmatrix} D_S & -\beta^* \\ \beta & D_Q \end{pmatrix}$$

with respect to the isomorphism (3.4), where  $\beta \in \mathcal{E}_{1,0}(X, \text{Hom}(S, Q))$  is the second fundamental form, see [10]. We shall now modify the connection  $D = D_E$  to  $D_b = D - \gamma_b$ , where  $\gamma_b = D'_{\text{End}E} j j^*$ . It turns out that  $\gamma = g^* \circ \beta \circ j^*$ , thus  $\gamma \wedge \gamma = 0$ , and that  $D_{\text{End}E} \gamma = \bar{\partial} \gamma$ . Moreover, it follows that

$$D_b \sim \begin{pmatrix} D_S & * \\ 0 & D_Q \end{pmatrix}$$

and hence

$$(3.6) \quad \Theta_b \sim \begin{pmatrix} \Theta_S & * \\ 0 & \Theta_Q \end{pmatrix},$$

so that  $c(D_b) = c(D_S)c(D_Q)$ . If  $D_t = D - t\gamma_b$  we have  $\Theta_t = \Theta - t\bar{\partial}\gamma_b$ ; thus it has bidegree  $(1, 1)$ . If we let

$$(3.7) \quad b = \int_0^1 \int_e \aleph \tilde{\gamma}_b \wedge e^{\tilde{I} + \aleph \tilde{\Theta} - t \aleph \bar{\partial} \tilde{\gamma}_b} = \sum_{\ell \geq 0} \int_e \aleph \tilde{\gamma}_b \wedge e^{\tilde{I} + \aleph \tilde{\Theta}} \wedge \frac{1}{(\ell + 1)!} (-\aleph \bar{\partial} \tilde{\gamma}_b)^\ell$$

it follows from (2.6) that  $db = c(D_E) - c(D_S)c(D_Q)$ . Moreover, if  $L = j j^* / (1 - t)$ , then (3.1) holds. In fact,  $\dot{D} = -\gamma_b$ , and  $[j j^*, g^* \circ \beta \circ j^*] = g^* \circ \beta \circ j^*$ , so that

$$(3.8) \quad D'_{\text{End}E, t} L = D'_{\text{End}E} L - t[\gamma_b, L] = \frac{1}{1 - t} \gamma_b - \frac{t}{1 - t} \gamma_b = \gamma_b.$$

**Proposition 3.2.** *If*

$$(3.9) \quad v = \sum_{\ell=1}^{m-1} \frac{(-1)^\ell}{2\ell} \int_e \widetilde{jj^*} \wedge (\tilde{I} + \mathfrak{N}\tilde{\Theta} - \mathfrak{N}\bar{\partial}\tilde{\gamma}_b)_{m-\ell-1} \wedge (-\mathfrak{N}\bar{\partial}\tilde{\gamma}_b)_\ell,$$

then  $-2\mathfrak{N}\partial v = b$ .

*Proof.* Observe that

$$\partial \int_0^{1-\epsilon} \int_e \frac{\widetilde{jj^*}}{1-t} \wedge e^{\tilde{I} + \mathfrak{N}\Theta_1} dt = \int_0^{1-\epsilon} \int_e \frac{D_1 \widetilde{jj^*}}{1-t} \wedge e^{\tilde{I} + \mathfrak{N}\Theta_1} dt = 0,$$

since  $D_1 \widetilde{jj^*} = \widetilde{D_{\text{End}E,1} jj^*} = 0$  in view of Lemma 2.1 and (3.8). Therefore,

$$\mathfrak{N}\partial \int_0^{1-\epsilon} \int_e \widetilde{jj^*} \wedge \frac{e^{\tilde{I} + \mathfrak{N}\tilde{\Theta} - t\mathfrak{N}\bar{\partial}\tilde{\gamma}_b} - e^{\tilde{I} + \mathfrak{N}\tilde{\Theta} - \mathfrak{N}\bar{\partial}\tilde{\gamma}_b}}{1-t} dt = \int_0^{1-\epsilon} \int_e \mathfrak{N}\tilde{\gamma}_b \wedge e^{\tilde{I} + \mathfrak{N}\tilde{\Theta} - t\mathfrak{N}\bar{\partial}\tilde{\gamma}_b}.$$

The proposition now follows by letting  $\epsilon \rightarrow 0$  and computing the  $t$ -integral on the left hand side.  $\square$

Altogether we therefore have that  $-dd^c v = c(D_E) - c(D_S)c(D_Q)$  and thus  $\hat{c}(E) = \hat{c}(S)\hat{c}(Q)$ .

#### 4. PROOF OF THE MAIN FORMULA

Let  $f$  be a nontrivial holomorphic section of  $E$ ,  $Z = \{f = 0\}$ , and let  $S$  be the trivial subbundle of  $E$  over  $X \setminus Z$ , generated by the  $f$ . We then have the short exact sequence (3.3) over  $X \setminus Z$ , where  $g: E \rightarrow Q = E/Q$  is the natural projection. Let  $\sigma$  be the section of the dual bundle  $E^*$  with minimal norm such  $\sigma \cdot f = 1$ . Then clearly

$$(4.1) \quad \widetilde{jj^*} = f \wedge \sigma.$$

Observe that the natural conjugate-linear isometry  $E \simeq E^*$ ,  $\eta \mapsto \eta^*$ , defined by

$$\eta^* \cdot \xi = \langle \xi, \eta \rangle, \quad \xi \in \mathcal{E}(X, E),$$

extends to an isometry on the space of form-valued sections.

**Lemma 4.1.** *If  $\phi = -\partial \log |f|^2$ , then  $D'\sigma = \phi \wedge \sigma$ .*

*Proof.* Observe that  $\sigma = f^*/|f|^2$ . Since  $D = D_E$  is the Chern connection,  $D'f^* = (\bar{\partial}f)^* = 0$ , so we have

$$D'\sigma = D'(f^*/|f|^2) = \partial \frac{1}{|f|^2} \wedge f^* = -\partial \log |f|^2 \wedge \sigma.$$

$\square$

Following Section 3 we let  $\gamma_b = D'_{\text{End}E}(jj^*)$ . By Lemma 4.1 and (4.1) we then have

$$(4.2) \quad \tilde{\gamma}_b = (Df - f \wedge \phi) \wedge \sigma$$



and

$$(4.3) \quad \bar{\partial}\tilde{\gamma}_b = (Df - f \wedge \phi) \wedge \bar{\partial}\sigma + (\Theta f + f \wedge \bar{\partial}\phi) \wedge \sigma.$$

The following formula is the key point in the analysis of the singularities of  $c(D_Q)$ .

**Proposition 4.2.** *In  $X \setminus Z$  we have the explicit formula*

$$(4.4) \quad c(D_Q) = \int_e f \wedge \sigma \wedge e^{\tilde{I} + \aleph \tilde{\Theta} - \aleph Df \wedge \bar{\partial}\sigma}.$$

*Proof.* Since  $\Theta_b = \Theta - \bar{\partial}\gamma_b$  we have by (4.3) that

$$\tilde{\Theta}_b = \tilde{\Theta} - ((Df - f \wedge \phi) \wedge \bar{\partial}\sigma + (\Theta f + f \wedge \bar{\partial}\phi) \wedge \sigma).$$

For any section  $A$  of  $\text{End}(E)$ ,

$$(4.5) \quad \int_e f \wedge \sigma \wedge \tilde{A}_{m-1} = \int_e f \wedge \sigma \wedge e^{\tilde{A}}$$

is the determinant of the restriction of  $A$  to  $Q$ , that is, the determinant of  $gAg^*$ . In view of (3.6) therefore the expression on the right hand side of (4.4) is equal to  $\det(I_Q + \aleph \Theta_Q) = c(D_Q)$ .  $\square$

Now, let  $v$  and  $b$  be the forms in  $X \setminus Z$  defined by (3.7) and (3.9).

**Proposition 4.3.** (i) *The forms  $v$ ,  $b$ ,  $c(D_Q)$ , and  $c(D_S) \wedge c(D_Q)$  are locally integrable in  $X$ .*

(ii) *If the natural extensions are denoted by capitals, then*

$$(4.6) \quad -2\aleph \partial V = B,$$

and

$$(4.7) \quad -dd^c V = c(D_E) - C(D_S)C(D_Q).$$

*Proof.* This is clearly a local question at  $Z$ . Locally we can write  $f = f_1 e_j + \dots + f_m e_m$ , where  $e_j$  is a local holomorphic frame for  $E$ . In a small neighborhood  $U$  of a given point in  $X$ , Hironaka's theorem provides an  $n$ -dimensional complex manifold  $\tilde{U}$  and a proper mapping  $\Pi: \tilde{U} \rightarrow U$  which is a biholomorphism outside  $\Pi^{-1}(\{f_1 \dots f_\nu = 0\})$ , and such that locally on  $\tilde{U}$  there are holomorphic coordinates  $\tau$  such that  $\Pi^* f_j = u^j \tau_1^{\alpha-1} \dots \tau_n^{\alpha_n}$ , where  $u_j$  nonvanishing; i.e., roughly speaking  $\Pi^* f_j$  are monomials. By a resolution over a suitable toric manifold, following [3] and [18], we may assume in the same way that one of the functions so obtained divides the other ones. For simplicity we will make a slight abuse of notation and suppress all occurring  $\Pi^*$  and thus denote these functions by  $f_j$  as well. We may therefore assume that  $f = f_0 f'$  where  $f_0$  is a holomorphic function and  $f'$  is a non-vanishing section. Since  $\sigma = f^*/|f|^2$ , it follows that  $\sigma = \sigma'/f_0$  where  $\sigma'$  is smooth, and hence

$$\widetilde{jj^*} = f \wedge \sigma = f' \wedge \sigma'$$

is smooth in this resolution. Moreover,  $Df \wedge \bar{\partial}\sigma = Df' \wedge \bar{\partial}\sigma' + \dots$ , where  $\dots$  denote terms that contain some factor  $f'$  or  $\sigma'$ . In view of Proposition 4.2 it follows that (the pullback of)  $c(D_Q)$  is smooth, and therefore locally integrable. Since the push-forward of a locally integrable form is locally integrable we can conclude that  $c(D_Q)$  is locally integrable.

It follows that also  $\tilde{\gamma}_b = D'(f \wedge \sigma)$  and  $\bar{\partial}\tilde{\gamma}_b$  are smooth. Since (4.6) and (4.7) hold in  $X \setminus Z$  and  $c(D_E)$  is smooth, it follows that all the forms are smooth in the resolution. We can conclude that all the forms are locally integrable in  $X$  and that (4.6) and (4.7) hold.  $\square$

The presence of the factor  $\widetilde{jj^*} = f \wedge \sigma$  implies that, cf., (3.9),

$$(4.8) \quad v = \sum_{\ell=1}^{m-1} \frac{(-1)^\ell}{2^\ell} \int_e f \wedge \sigma \wedge (\tilde{I} + \mathfrak{s}\tilde{\Theta} - \mathfrak{s}Df \wedge \bar{\partial}\sigma)_{m-1-\ell} \wedge (-\mathfrak{s}Df \wedge \bar{\partial}\sigma)_\ell.$$

**Definition 1.** We define the current  $W$  as

$$(4.9) \quad W = \log(1/|f|)c(D_Q) - V = \\ \log(1/|f|) \int_e f \wedge \sigma \wedge (\mathfrak{s}\tilde{\Theta} + \tilde{I} - \mathfrak{s}Df \wedge \bar{\partial}\sigma)_{m-1} \\ - \sum_{\ell=1}^{m-1} \frac{(-1)^\ell}{2^\ell} \int_e f \wedge \sigma \wedge (\tilde{I} + \mathfrak{s}\tilde{\Theta} - \mathfrak{s}Df \wedge \bar{\partial}\sigma)_{m-1-\ell} \wedge (-\mathfrak{s}Df \wedge \bar{\partial}\sigma)_\ell.$$

In particular, if  $E$  is a line bundle, i.e.,  $m = 1$ , then  $V = 0$ , and since  $\sigma \cdot f = 1$  we have that  $W = \log(1/|f|)$ . It is now a simple matter to conclude the proof of Theorem 1.1.

*Proof of Theorem 1.1.* Consider a resolution of singularities in which  $f = f_0 f'$  with  $f'$  non-vanishing, as in the proof of Proposition 4.3. Then we know that  $c(D_Q)$  is smooth, and therefore  $\log|f|c(D_Q)$  is locally integrable there. Moreover, since  $\log|f| = \log|f_0| + \log|f'|$  we have that

$$\lambda|f|^{2\lambda} \frac{\mathfrak{s}\partial|f|^2 \wedge \bar{\partial}|f|^2}{|f|^4} \wedge c(D_Q) = \\ \lambda|f_0|^{2\lambda}|f'|^{2\lambda} \mathfrak{s} \left( \frac{df_0}{f_0} + \frac{\partial|f'|^2}{|f'|^2} \right) \wedge \left( \frac{d\bar{f}_0}{\bar{f}_0} + \frac{\bar{\partial}|f'|^2}{|f'|^2} \right) \wedge c(D_Q).$$

This form is locally integrable for  $\lambda > 0$  and tends to

$$[f_0 = 0] \wedge c(D_Q) = dd^c(\log|f|c(D_Q)) \mathbf{1}_{\{f_0=0\}}$$

when  $\lambda \rightarrow 0$ , where  $[f_0 = 0]$  is the current of integration over the divisor defined by  $f_0$ . Thus  $M$  is a closed current of bidegree  $(*, *)$  and order zero in  $X$  with support on  $Z$ . Thus, see, e.g., [10],  $M_k = 0$  for  $k < p = \text{codim } Z$  and  $M_p = \sum_j \alpha_j Z_j^p$  for some numbers  $\alpha_j$ . To see that

$\alpha_j$  is precisely the multiplicity of  $f$  on  $Z_j^p$  we can locally deform the Hermitian metric to a trivial metric. Then  $\Theta = 0$  and a straight-forward computation, see [2], reveals that  $c_{p-1}(D_Q) = (dd^c \log |f|)^{p-1}$ . Therefore,  $M = dd^c(\log |f|(dd^c \log |f|)^{p-1})$  which is equal to the multiplicity times  $[Z_j^p]$  according to King's formula, see [11] and [10]. Thus part (i) of the theorem is proved. Since  $c(D_S) - 1 = c_1(D_S) = dd^c \log(1/|f|)$  we have

$$dd^c(\log(1/|f|)c(D_Q)) = C(D_S) \wedge C(D_Q) - C(D_Q) - dd^c(\log |f|c(D_Q))\mathbf{1}_Z.$$

Now part (ii) follows from Proposition 4.3, cf, (4.9).  $\square$

## 5. A DIRECT APPROACH TO (1.6)

We use the same notation as in the previous section. In [6], Berndtsson introduced the deformation  $D_a = D - \gamma_a$  of  $D$  on  $E$ , where

$$(5.1) \quad \tilde{\gamma}_a = Df \wedge \sigma,$$

in order to construct Koppelman formulas for  $\bar{\partial}$  on manifolds. He proved formula (5.7) below for  $k = m$  (i.e.,  $\bar{\partial}a_m = da_m = c_m(E)$ ). For the general case first we must understand the geometric meaning of  $D_a$ . Since  $D_af = 0$ , we have that  $D_a\xi$  is in  $S$  if  $\xi$  is a section of  $S$ . Moreover, if  $\xi$  is a section of  $S^\perp$ , then  $D_a\xi = D_E\xi$ . Now

$$(5.2) \quad g\xi \mapsto g(D_a\xi)$$

is a well-defined connection on  $Q$ , and we claim that it is actually the Chern connection  $D_Q$ . In fact, if  $\eta = g\xi$ , then

$$D_Q\eta = g(D_E(g^*\eta)) = g(D_a(g^*\eta)) = g(D_a\xi).$$

It follows that  $\Theta_Q\eta = g(\Theta_a\xi)$ , and since  $\Theta_a\xi = 0$  if  $\xi$  takes values in  $S$ , we have that

$$(5.3) \quad \mathbb{N}\Theta_a \sim \begin{pmatrix} 0 & * \\ 0 & \mathbb{N}\Theta_Q \end{pmatrix}$$

with respect to the smooth isomorphism (3.4). Therefore,

$$\mathbb{N}\Theta_a + I_E \sim \begin{pmatrix} I_S & * \\ 0 & I_Q + \mathbb{N}\Theta_Q \end{pmatrix},$$

and taking the determinant, we find that

$$(5.4) \quad c(D_Q) = c(D_a).$$

**Proposition 5.1.** *If  $\gamma_a$  is defined by (5.1), then*

$$(5.5) \quad -tD\tilde{\gamma}_a + t^2\widetilde{\gamma_a \wedge \gamma_a} = -t(Df \wedge \bar{\partial}\sigma + \Theta f \wedge \sigma) + (t - t^2)Df \wedge \phi \wedge \sigma.$$

*Proof.* A simple computation yields

$$D\tilde{\gamma}_a = \Theta f \wedge \sigma + Df \wedge \bar{\partial}\sigma + Df \wedge \phi \wedge \sigma$$

and

$$\widetilde{\gamma_a \wedge \gamma_a} = Df \wedge \sigma \cdot Df \wedge \sigma,$$

where the dot means the natural contraction of  $E$  and  $E^*$  so that  $\xi \cdot (\alpha \wedge \eta) = \alpha(\xi \cdot \eta)$  if  $\xi$  and  $\eta$  are sections of  $E$  and  $E^*$ , respectively, and  $\alpha$  is a form. Since  $\sigma \cdot Df = -D'\sigma \cdot f = \phi$  we get the desired formula.  $\square$

**Proposition 5.2.** *If*

$$(5.6) \quad a = \int_e \mathbb{N} Df \wedge \sigma \wedge e^{\tilde{I} + \mathbb{N}\tilde{\Theta}} \wedge \sum_{\ell=0}^{\infty} \frac{(-\mathbb{N} Df \wedge \bar{\partial}\sigma)^\ell}{(\ell+1)!}$$

then

$$(5.7) \quad \bar{\partial}a = da = c(D_E) - c(D_Q)$$

in  $X \setminus Z$ .

*Proof.* We choose the homotopy  $D_t = D - t\gamma_a$  between  $D = D_0$  and  $D_1 = D_a$ . In view of (2.6), (2.1), and Proposition 5.1 we have that

$$a = \int_e \int_0^1 \mathbb{N} Df \wedge \sigma \wedge e^{\tilde{I} + \mathbb{N}\tilde{\Theta} - t\mathbb{N}(\Theta f \wedge \sigma + Df \wedge \bar{\partial}\sigma) - (t-t^2)Df \wedge \phi \wedge \sigma} dt$$

satisfies the second equality in (5.7) in  $X \setminus Z$ . Noticing that  $\sigma \wedge \sigma = 0$ , a computation of the  $t$ -integral yields (5.6). Since  $a$  has bidegree  $(*, * - 1)$  and  $da$  has bidegree  $(*, *)$  it follows that  $\bar{\partial}a = da$ .  $\square$

The forms  $a$  and  $b$  are related in the following way.

**Proposition 5.3.** *In  $X \setminus Z$  we have that*

$$(5.8) \quad b = a + \mathbb{N} \partial \log |f|^2 \wedge c(D_Q)$$

*Proof.* Starting with (3.7) we have

$$\begin{aligned} b &= \int_e \mathbb{N}(Df - f \wedge \phi) \wedge \sigma \wedge e^{\tilde{I} + \mathbb{N}\tilde{\Theta}} \wedge \sum_{\ell=0}^{\infty} \frac{(-\mathbb{N} Df + \mathbb{N} f \wedge \phi)^\ell}{(1+\ell)!} \wedge (\bar{\partial}\sigma)^\ell = \\ &\quad - \int_e e^{\tilde{I} + \mathbb{N}\tilde{\Theta}} \wedge \sum_{\ell=0}^{\infty} \frac{(-\mathbb{N} Df + \mathbb{N} f \wedge \phi)^{\ell+1}}{(\ell+1)!} \wedge \sigma \wedge (\bar{\partial}\sigma)^\ell = \\ &\quad - \int_e e^{\tilde{I} + \mathbb{N}\tilde{\Theta} - \mathbb{N} Df + \mathbb{N} f \wedge \phi} \wedge \sum_{\ell=0}^{\infty} \sigma \wedge (\bar{\partial}\sigma)^\ell = \\ &\quad - \int_e e^{\tilde{I} + \mathbb{N}\tilde{\Theta} - \mathbb{N} Df} \wedge (1 + \mathbb{N} f \wedge \phi) \wedge \sum_{\ell} \sigma \wedge (\bar{\partial}\sigma)^\ell. \end{aligned}$$

In view of (6.3) and (6.6), recalling that  $\phi = -\partial \log |f|^2$ , we now get (5.8).  $\square$

By a resolution of singularities as in the proof of Proposition 4.3 above one can see that  $a$  is locally integrable. Let  $A$  denote its natural extension. By such a resolution one can also verify that the formal

computation (using Proposition 5.3)  $-2\aleph\partial(\log(1/|f|)c(D_Q) - V) = B - \aleph\partial\log|f|^2\wedge C(D_Q) = A$  is legitimate, and thus we have

$$(5.9) \quad A = -2\aleph\partial W.$$

As a consequence we get that  $\bar{\partial}A = dA = c(D_E) - c(D_Q) - M$ .

## 6. FACTORIZATION OF CURRENTS

Since  $a$  and  $c(D_Q)$  are locally integrable,  $|f|^{2\lambda}a$  and  $|f|^{2\lambda}c(D_Q)$  are well-defined currents for  $\operatorname{Re} \lambda > -\epsilon$  and we have

$$(6.1) \quad A = |f|^{2\lambda}a|_{\lambda=0} \quad \text{and} \quad C(D_Q) = |f|^{2\lambda}c(D_Q)|_{\lambda=0}.$$

It also follows that

$$(6.2) \quad M = -d|f|^{2\lambda}\wedge a|_{\lambda=0} = -\bar{\partial}|f|^{2\lambda}\wedge a|_{\lambda=0}.$$

Now consider the expression (5.6) for  $a$ . Since each term in  $\exp(\tilde{I} + \aleph\tilde{\Theta})$  has the same degree in  $e_j$  and  $e_k^*$  it must be multiplied by terms with the same property in order to get a product with full degree. Therefore we can rewrite  $a$  as

$$(6.3) \quad a = - \int_e e^{\tilde{I} + \aleph\tilde{\Theta} - \aleph Df} \wedge \sum_0^\infty \sigma \wedge (\bar{\partial}\sigma)^\ell.$$

In [1] we introduced the currents

$$U = |f|^{2\lambda} \frac{\sigma}{1 - \bar{\partial}\sigma} \Big|_{\lambda=0} = |f|^{2\lambda} \wedge \sigma \wedge \sum_\ell (\bar{\partial}\sigma)^{\ell-1} \Big|_{\lambda=0}$$

and

$$R = \bar{\partial}|f|^{2\lambda} \wedge \frac{\sigma}{1 - \bar{\partial}\sigma} \Big|_{\lambda=0} = \bar{\partial}|f|^{2\lambda} \wedge \sigma \wedge \sum_\ell (\bar{\partial}\sigma)^{\ell-1} \Big|_{\lambda=0}.$$

It is part of the statement that the right hand sides are current valued holomorphic functions for  $\lambda > -\epsilon$ , evaluated at  $\lambda = 0$ . In general  $U$  and  $R$  are *not* locally integrable. The current  $R$  is supported on  $Z$ ,

$$R = R_p + \cdots + R_{\min(m,n)},$$

where  $R_k$  is the component of bidegree  $(0, k)$  taking values in  $\Lambda^k E^*$ , and  $(\delta_f - \bar{\partial})U = 1 - R$ . In view of (6.3), (6.1), and (6.2) we have the factorization formulas

$$(6.4) \quad M = \int_e e^{\aleph\tilde{\Theta} + \tilde{I} - \aleph Df} \wedge R,$$

$$(6.5) \quad A = - \int_e e^{\aleph\tilde{\Theta} + \tilde{I} - \aleph Df} \wedge U,$$

and moreover, cf. (4.4),

$$(6.6) \quad C(D_Q) = \int_e f \wedge \sigma \wedge e^{\aleph\tilde{\Theta} + \tilde{I} - \aleph Df} \wedge \bar{\partial}\sigma = \int_e e^{\aleph\tilde{\Theta} + \tilde{I} - \aleph Df} \wedge f \wedge U.$$

## 7. POSITIVITY

Let  $E \rightarrow X$  be a Hermitian holomorphic bundle as before and let  $e_j$  be an orthonormal local frame. A section

$$A = i \sum_{jk} A_{jk} \otimes e_j \otimes e_k^*$$

of  $T_{1,1}^*(X) \otimes \text{End}(E)$  is Hermitian if  $A_{jk} = -\overline{A_{kj}}$ . It then induces a Hermitian form  $a$  on  $T^{1,0}(X) \otimes E^*$  by

$$a(\xi \otimes e_j^*, \eta \otimes e_k^*) = A_{jk}(\xi, \bar{\eta}),$$

if  $\xi, \eta$  are  $(1,0)$ -vectors. We say that  $A$  is (Bott-Chern) positive,  $A \geq_B 0$  if the form  $a$  is positively semi-definite. In the same way any Hermitian  $A$  induces a Hermitian form  $a'$  on  $T^{1,0}(X) \otimes E$  and it is called Nakano positive,  $A \geq_N 0$ , if  $a'$  is positively semi-definite.

Notice that  $\kappa\Theta$  is Hermitian; it is said to be Nakano positive if  $\kappa\Theta \geq_N 0$ . Analogously we say that  $E$  is positive,  $E \geq_B 0$ , if  $\kappa\Theta \geq_B 0$ . Neither of these positivity concepts implies the other one unless  $m = 1$ .

Since  $\Theta_{jk}(E^*) = -\Theta_{jk}(E)$  it follows that  $E$  is positive in our sense if and only if  $E^*$  is Nakano negative. The next proposition explains the interest of Bott-Chern positivity in this context.

**Proposition 7.1.** *Let*

$$(7.1) \quad 0 \rightarrow S \rightarrow E \rightarrow Q \rightarrow 0$$

*be a short exact sequence of Hermitian holomorphic vector bundles. Then  $E \geq_B 0$  implies that  $Q \geq_B 0$ .*

*Proof.* It is well-known, see for instance [10], that  $E \leq_N 0$  implies that  $S \leq_N 0$ . From the sequence (7.1) above we get the exact sequence  $0 \rightarrow Q^* \rightarrow E^* \rightarrow S^* \rightarrow 0$ . Since  $E^* \leq_N 0$  implies  $Q^* \leq_N 0$ , it follows that  $E \geq_B 0$  implies  $Q \geq_B 0$ .  $\square$

The next simple lemma reveals that our definition of Bott-Chern positivity coincides with the one used in [8].

**Lemma 7.2.**  *$A \geq_B 0$  if and only if there are sections  $f_\ell$  of  $T_{1,0}^*(X) \otimes E$  such that*

$$(7.2) \quad A = i \sum_{\ell} f_\ell \otimes f_\ell^*.$$

Observe that if  $f_\ell = \sum f_j^\ell \otimes e_j$ , then  $f_\ell^* = \sum \bar{f}_j^\ell \otimes e_j^*$  since  $e_j$  is orthonormal.

*Proof.* If (7.2) holds, then

$$a(\xi, \xi) = \sum_{\ell} f_\ell(\xi) f_\ell^*(\xi^*) = \sum_{\ell} |f_\ell(\xi)|^2 \geq 0$$

for all  $\xi$  in  $T^{1,0} \otimes E^*$ . Conversely, if  $a$  is positive, it is diagonalizable, and so there is a basis  $f_\ell$  for  $T_{1,0}^* \otimes E$  such that (7.2) holds.  $\square$

If we identify  $f_\ell$  with  $\sum f_j^\ell \wedge e_j$  as before, then (7.2) means that

$$(7.3) \quad \tilde{A} = -i \sum_{\ell} f_\ell \wedge f_\ell^*.$$

If  $B = \sum B_{jk} e_j \otimes e_k^*$  is a scalar-valued section of  $\text{End} E$ , then it is Hermitian if and only if  $B_{jk} = \bar{B}_{kj}$  and it is positively semi-definite if and only if

$$B = \sum_{\ell} g_\ell \otimes g_\ell^*$$

for some sections  $g_\ell$  of  $E$ ; or equivalently,

$$(7.4) \quad \tilde{B} = \sum_{\ell} g_\ell \wedge g_\ell^*.$$

**Proposition 7.3.** *Assume that  $A_j$  are  $(1,1)$ -form-valued Hermitian sections of  $E$  and  $B_k$  scalarvalued sections, such that  $A_j \geq_B 0$  and  $B_k \geq 0$ . Then*

$$(7.5) \quad \int_e \tilde{A}_1 \wedge \dots \wedge \tilde{A}_r \wedge \tilde{B}_{r+1} \wedge \dots \wedge \tilde{B}_m$$

is a positive  $(r, r)$ -form.

*Proof.* In view of (7.3) and (7.4), we see that (7.5) is a sum of terms like

$$\begin{aligned} & \int_e (-i)^r f_1 \wedge f_1^* \wedge \dots \wedge f_r \wedge f_r^* \wedge g_{r+1} \wedge g_{r+1}^* \wedge \dots \wedge g_m \wedge g_m^* = \\ & (-i)^r c_{m-r} \int_e f_1 \wedge \dots \wedge f_r \wedge \dots \wedge g_m \wedge f_1^* \wedge \dots \wedge f_r^* \wedge \dots \wedge g_m^* = \\ & (-i)^r c_{m-r} \int_e \omega \wedge e_1 \wedge \dots \wedge e_m \wedge \bar{\omega} \wedge e_1^* \wedge \dots \wedge e_m^*, \end{aligned}$$

where  $\omega$  is an  $(r, 0)$ -form and  $c_p = (-1)^{p(p-1)/2} = i^{p(p-1)}$ . By further simple computations,

$$\begin{aligned} & (-i)^r c_{m-r} (-1)^{mr} \int_e \omega \wedge \bar{\omega} \wedge e_1 \wedge \dots \wedge e_m \wedge e_1^* \wedge \dots \wedge e_m^* = \\ & (-i)^r c_{m-r} (-1)^{mr} c_m \omega \wedge \bar{\omega} = i^{r^2} \omega \wedge \bar{\omega} \end{aligned}$$

the proposition follows, since the last form is positive.  $\square$

**Proposition 7.4.** *If  $E \geq_B 0$  (or  $E \geq_N 0$ ), then the Chern forms  $c_k(D_E)$  are positive for all  $k$ .*

*Proof.* Since  $\alpha\Theta \geq_B 0$  by assumption, and clearly  $I \geq 0$ , it follows from Proposition 7.3 that

$$c_k(D_E) = \int_e (\mathfrak{s}\tilde{\Theta})_k \wedge \tilde{I}_{m-k}$$

is positive.  $\square$

*Proof of Theorem 1.2.* We have just seen that  $c(D_E) \geq 0$ . From (1.3) it follows that the current  $M_k$  is positive if  $c_{k-1}(D_Q)$  is positive. From (4.4) we have that

$$(7.6) \quad c_{k-1}(D_Q) = \int_e f \wedge \sigma \wedge (\mathfrak{s}\tilde{\Theta} - \mathfrak{s}Df \wedge \bar{\partial}\sigma)_{k-1} \wedge \tilde{I}_{m-k} = \sum_{j=1}^{k-1} \int_e f \wedge \sigma \wedge (\mathfrak{s}\tilde{\Theta})_{k-1-j} \wedge (-\mathfrak{s}Df \wedge \bar{\partial}\sigma)_j \wedge \tilde{I}_{m-k}.$$

If  $s = f^*$  as before, then  $\sigma = s/|f|^2$ , and therefore we have

$$(7.7) \quad c_{k-1}(D_Q) = \sum_{j=1}^{k-1} \int_e \frac{f \wedge s}{|f|^2} \wedge \left( \frac{-\mathfrak{s}Df \wedge \bar{\partial}s}{|f|^2} \right)_j \wedge (\mathfrak{s}\tilde{\Theta})_{k-1-j} \wedge \tilde{I}_{m-k}.$$

Since  $\bar{\partial}s = (Df)^*$  it now follows immediately from Proposition 7.3 that  $c_k(D_Q)$  is positive if  $\mathfrak{s}\tilde{\Theta} \geq_B 0$ .

It remains to see that one can choose  $W$  so that it is positive where  $|f| < 1$ . Notice that if some of the  $A_j$  in (7.5) are replaced by  $A'_j \geq_B A_j$ , then the resulting form will be larger; this follows immediately from the proof. Now,  $\log(1/|f|)c(D_Q)$  is positive when  $|f| < 1$ . From (4.8) we have that

$$v_k = \sum_{\ell=1}^k \frac{(-1)^\ell}{2\ell} \int_e f \wedge \sigma \wedge (\mathfrak{s}\tilde{\Theta} - \mathfrak{s}Df \wedge \bar{\partial}\sigma)_{k-\ell} \wedge (-\mathfrak{s}Df \wedge \bar{\partial}\sigma)_\ell \wedge \tilde{I}_{m-k-1}.$$

Since this is an alternating sum of positive terms it has no sign. If we replace each factor  $-\mathfrak{s}Df \wedge \bar{\partial}\sigma$  by  $\mathfrak{s}\tilde{\Theta} - \mathfrak{s}Df \wedge \bar{\partial}\sigma$ , then we get a larger form which in addition is closed, since it is just a certain constant times  $c_k(D_Q)$ , cf., (7.6). Therefore, for a suitable constant  $\nu_k$   $-v'_k = -v_k + \nu_k c_k(D_Q)$  is a positive form and  $dv'_k = dv_k$ . Thus the current

$$W'_k = -V_k + \nu_k C_k(D_Q) + \log(1/|f|)C_k(D_Q)$$

will have the stated property.  $\square$

The modification of  $v$  in last part of the proof is precisely as in [8] but with our notation, and for an arbitrary  $k$  rather than just  $k = m - 1$ . It is not necessary to consider each  $v_k$  separately. By the same argument one can see directly that  $-v' = -v + \nu c(D_Q)$  is positive if  $\nu$  is appropriately chosen, and  $dv' = dv$ .

One can prove that if we multiply (7.7) with  $\lambda \partial|f|^2 \wedge \bar{\partial}|f|^2 / |f|^2$  and let  $\lambda \rightarrow 0^+$ , then all terms with  $j < p - 1$  will disappear; see for instance the proof of Theorem 1.1 in [1]. We thus have



**Proposition 7.5.** *If  $p = \text{codim} \{f = 0\}$ , then*

$$M_k = \lim_{\lambda \rightarrow 0^+} \lambda |f|^{2\lambda} \frac{\partial |f|^2 \wedge \bar{\partial} |f|^2}{|f|^2} \wedge \sum_{j=p-1}^{k-1} \int_e \frac{f \wedge s}{|f|^2} \wedge \left( \frac{-\mathfrak{s} D f \wedge \bar{\partial} s}{|f|^2} \right)_j \wedge (\mathfrak{s} \tilde{\Theta})_{k-1-j} \wedge \tilde{I}_{m-k}.$$

From this formula it is apparant that  $M_k$  vanishes if  $k < p$ , and that  $M_p$  is positive, regardless of  $\mathfrak{s}\Theta$ . One can also derive this formula from (6.4).

*Remark 2.* When  $k > p$ ,  $M_k$  depends on the metric, but there is still a certain uniqueness: Let  $Z^k$  be the union of the irreducible components  $Z_j^k$  of  $Z$  of codimension  $k$ . One can verify, see [2], that the restriction of  $M_k$  to  $Z^k$  is a sum

$$\sum_j \alpha_j^k [Z_j^k],$$

where  $\alpha_j^k$  are nonnegative numbers that are independent of the metric. However the geometric meaning of these numbers is not clear to us.  $\square$

## 8. SOME EXAMPLES

The first two examples suggest that not only the component  $M_p$  of the current  $M$  is of interest.

*Example 1.* Let us assume that  $X$  is compact, and that we have sections  $f_j$  of rank  $m_j$  bundles  $E_j \rightarrow X$ , such that  $\sum m_j = n$ . If  $E = \oplus E_j$  and  $f = (f_1, \dots, f_r)$ , then the intersection number  $\nu$  of the varieties  $Z_j = \{f_j = 0\}$  is equal to the integral of

$$c_n(E) = c_{m_1}(E_1) \wedge \dots \wedge c_{m_r}(E_r)$$

over  $X$ . Since  $M_n$  represents the cohomology class  $c_n(E)$ , we thus get the representation

$$\nu = \int_X M_n,$$

i.e., an integral over the set-theoretic intersection  $Z = \cap Z_j$ . If  $E$  is positive then  $M_n$  is positive. If  $Z$  is discrete, i.e.,  $f$  is a complete intersection, then  $M_n = [Z]$ , and in this case thus we just get the sum of the points in  $Z$  counted with multiplicities, as expected.  $\square$

*Example 2.* Let  $X$  be a compact Kähler manifold with metric form  $\omega$ , and let  $f$  be a holomorphic section of  $E \rightarrow X$ . If moreover  $E \geq_B 0$ , then we know that  $c(D_E)$ ,  $M$ , and  $c(D_Q)0$  are all positive. Because of (1.4), we therefore have that

$$\int_X M_k \wedge \omega_{n-k} = \int_X c_k(D_E) \wedge \omega_{n-k} - \int_X c_k(D_Q) \wedge \omega_{n-k} \leq \int_X c_k(D_E) \wedge \omega_{n-k}.$$

Thus we get an upper bound of the total mass of  $M_k$  in terms of the Chern class  $c_k(E)$ . Taking  $k = p = \text{codim } Z$  we get the estimate

$$\text{area}(Z^p) = \int_X [Z^p] \leq \int_X c_p(E) \wedge \omega_{n-p}.$$

□

*Example 3.* Now assume that  $X = \mathbb{P}^n$ , let

$$\omega = \ast \partial \bar{\partial} \log |z|^2 = dd^c \log |z|$$

denote the Fubini-Study metric and notice that

$$\int_{\mathbb{P}^n} \omega^n = 1,$$

that is, the total area of  $\mathbb{P}^n$  is  $1/n!$ .

Assume that  $F_1, \dots, F_m$  are polynomials in  $\mathbb{C}^n$  which form a complete intersection. If  $F_j$  has degree  $d_j$  (depending on  $z' = (z'_1, \dots, z'_n)$ ) then the homogenization  $f_j(z) = z_0^{d_j} F_j(z'/z_0)$  is a  $d_j$ -homogeneous polynomial in  $\mathbb{C}^{n+1}$  and hence corresponds to a section of the line bundle  $\mathcal{O}(d_j) \rightarrow \mathbb{P}^n$ . Thus  $f = (f_1, \dots, f_m)$  is a section of  $E = \oplus \mathcal{O}(d_j)$ . If  $E$  is equipped with the natural metric, i.e.,

$$\|h([z])\|^2 = \sum_j \frac{|h_j(z)|^2}{|z|^{2d_j}}$$

for a section  $h = \oplus h_j$  of  $E$  (here  $[z]$  denotes the point on  $\mathbb{P}^n$  corresponding to the point  $z \in \mathbb{C}^{n+1} \setminus \{0\}$  under the usual projection), then it is easy to check that  $E \geq_B 0$ . Therefore  $M_m \geq 0$ , and since moreover,

$$M_m|_{\mathbb{C}^n} = [Z],$$

if  $Z$  here denotes the zero variety  $\{F = 0\}$  in  $\mathbb{C}^n$ , then

$$\text{area}(Z) = \int_{\mathbb{C}^n} [Z] \wedge \omega_{n-m} \leq \int_{\mathbb{P}^n} M_m \wedge \omega_{n-m} = \int_{\mathbb{P}^n} c_m(D_E) \wedge \omega_{n-m},$$

since  $c_m(D_Q) = 0$ . Here “area” refers to the projective area of course. However,  $c(D_E) = (1 + d_1\omega) \wedge \dots \wedge (1 + d_m\omega)$ , and so

$$c_m(D_E) = d_1 \cdots d_m \omega^m.$$

Hence

$$\text{area}(Z) \leq d_1 \cdots d_m \frac{1}{(n-m)!}.$$

We also notice that the deviation from equality is precisely the total mass of  $M_m$  on the hyperplane at infinity. If  $m = n$  we get Bezout’s theorem

$$\#\{F = 0\} \leq d_1 \cdots d_n.$$

□

*Example 4.* If  $f$  is a complete intersection, i.e.,  $p = m$ , and  $W_{m-1}$  denotes the component of bidegree  $(m-1, m-1)$ , then

$$dd^c W_{m-1} = c_m(D_E) - [Z];$$

this means that  $W_{m-1}$  is a Green current for the cycle  $Z = \sum \alpha_j Z_j$ .

In the case when  $E = L_1 \oplus \dots \oplus L_m$  for some line bundles  $L_k$ , hence  $c_m(D_E) = c_1(D_{L_1}) \wedge \dots \wedge c_m(D_{L_m})$ , and  $f = (f_1, \dots, f_m)$ , where  $f_j$  are holomorphic sections of  $L_j$ , such a Green current was obtained already in [3].  $\square$

*Example 5.* Let  $X$  be a compact manifold such that there is a holomorphic section  $\eta$  of some vector bundle  $H \rightarrow X \times X$  that defines the diagonal  $\Delta \subset X \times X$ ; for instance  $X$  can be complex projective space. From Theorem 1.1 we get a current  $W_n$  such that  $dd^c W = c_n(D_H) - [\Delta]$ . If we let  $K(\zeta, z) = -W_n$  and  $P(\zeta, z) = c_n(D_H)$ , then

$$dd^c K = [\Delta] - P,$$

and this leads to the Koppelman type formula

$$(8.1) \quad \phi(z) - \int P(\zeta, z) \wedge \phi(\zeta) = \\ dd^c \int_X K \wedge \phi - d \int_X K \wedge d^c \phi + d^c \int_X K \wedge d\phi + \int_X K \wedge dd^c \phi$$

for the  $dd^c$ -operator. In particular, if  $\phi$  is closed  $(k, k)$ -form such that  $d\phi = 0$ , then  $d^c \phi = 0$  as well, and thus

$$v = \int_X K \wedge \phi$$

is an explicit solution to  $dd^c v = \phi - \int P \wedge \phi$ . However if  $X$  is non-compact one gets boundary integrals. It would be desirable to refine the construction to include somehow an appropriate line bundle with a metric that vanishes at the boundary, in order to obtain  $dd^c$ -formulas for, say, domains in  $\mathbb{C}^n$ .  $\square$

*Example 6.* Assume that  $f$  is a holomorphic section of some Hermitian bundle  $E \rightarrow X$  with zero variety  $Z$ . If  $f$  is locally a complete intersection we have seen that the current  $W_{m-1}$  from Theorem 1.1 is a Green current for  $[Z]$ . In general we have that  $dd^c W_{p-1} = c_p(D_E) - c_p(D_Q) - [Z^p]$  so we only get a current  $w$  such that  $dd^c w = [Z^p] - \gamma$ , where  $\gamma$  is locally integrable. However, there is another and simpler way to find such a current  $w$ , due to Meo, [17].

*Proposition 8.1 (Meo).* *Let  $f$  be a holomorphic section of a Hermitian vector bundle  $E \rightarrow X$ . The forms*

$$w = \log |f| (dd^c \log |f|)^{p-1} \mathbf{1}_{X \setminus Z}$$

and

$$\gamma = -(dd^c \log |f|)^p \mathbf{1}_{X \setminus Z}$$

are locally integrable on  $X$  and

$$(8.2) \quad dd^c w = [Z^p] - \gamma.$$

For the reader's convenience we provide a simple proof based on Hironaka's theorem.

*Sketch of proof.* Let  $f = f_0 f'$  be as before, i.e.,  $f_0$  is holomorphic and  $f'$  is a non-vanishing section. Then  $\log |f| = \log |f_0| + \log |f'|$ , and hence  $dd^c \log |f'|$  is smooth and  $dd^c \log |f_0| = [f_0 = 0]$  has support on the inverse image  $\tilde{Z}$  of  $Z$  in the resolution. Thus

$$w = (\log |f_0| + \log |f'|)(dd^c \log |f'|)^{p-1}, \quad \gamma = (dd^c \log |f'|)^p$$

are both locally integrable in the resolution and hence also on the original manifold. Moreover,

$$dd^c w = [f_0 = 0] \wedge (dd^c \log |f'|)^{p-1} + \gamma,$$

in particular  $(dd^c w)\mathbf{1}_{\tilde{Z}}$  is closed, and hence  $T = (dd^c w)\mathbf{1}_Z$  is a closed current on  $X$  of order zero. Therefore  $T = \sum \alpha_j [Z_j^p]$ , and since we can deform the metric into a trivial metric locally, it follows from King's formula, [11], that  $\alpha_j$  are precisely the multiplicities of  $f$  on  $Z_j^p$ .  $\square$

Assume now that  $X$  is a compact manifold such that there exists a holomorphic section  $\eta$  of some Hermitian bundle  $H \rightarrow X \times X$  as in Example 5 above. If furthermore the kernel  $K$  is reasonably regular we can assume that

$$dd^c \int_{\zeta} K(\zeta, z) \wedge \psi(\zeta) = \psi(z) - \int_{\zeta} P(\zeta, z) \wedge \psi(\zeta)$$

for any  $(k, k)$ -current  $\psi$ . We then have the explicit solution

$$g = w + \int_{\zeta} K(\zeta, z) \wedge \gamma(\zeta)$$

to the Green equation  $dd^c g = [Z^p] - \alpha$ , where  $\alpha$  is the smooth form

$$\alpha = \int_{\zeta} P(\zeta, z) \wedge \gamma(\zeta).$$

$\square$

The last example will be elaborated in a forthcoming paper.

## REFERENCES

- [1] M. ANDERSSON: *Residue currents and ideals of holomorphic functions*, Bull. Sci. Math. **128** (2004), 481–512.
- [2] M. ANDERSSON: *Residue currents of holomorphic sections and Lelong currents*, Arkiv för matematik **43** (2005), 201–219.
- [3] C. BERENSTEIN & A. YGER: *Green currents and analytic continuation*, J. Anal. Math. **75** (1998), 1–50.
- [4] C. A. BERENSTEIN & R. GAY & A. VIDRAS & A. YGER: *Residue currents and Bezout identities*, Progress in Mathematics **114** Birkhäuser Verlag (1993).

- [5] C. BERENSTEIN & A. YGER: *Analytic residue theory in the non-complete intersection case*, J. Reine Angew. Math. **527** (2000), 203–235.
- [6] B. BERNDTSSON: *Cauchy–Leray forms and vector bundles*, Ann. Scient. Ec. Norm. Sup. **24** (1991), 319–337.
- [7] J.-M. BISMUT & H. GILLET & CH. SOULÉ: *Complex immersions and Araakelov geometry*, The Grothendieck Festschrift, Vol. I, 249–331, Progr. Math., 86, Birkhuser Boston, Boston, MA, (1990).
- [8] R. BOTT & S.-S. CHERN: *Hermitian vector bundles and the equidistribution of the zeroes of their holomorphic sections*, Acta Math. **114** (1965), 71–112.
- [9] S.-S. CHERN: *Transgression in associated bundles*, Internat. J. Math. **2** (1991), 383–393.
- [10] J-P DEMAILLY: *Complex Analytic and Differential Geometry*, Monograph Grenoble (1997).
- [11] J. R. KING: *A residue formula for complex subvarieties*, Proc. Carolina conf. on holomoprhic mappings and minimal surfaces, Univ. of North Carolina, Chapel Hill (1970), 43–56.
- [12] R. HARVEY & S. SEMMES: *Zero divisors of atomic functions*, Ann. of Math. **135** (1992), 567–600.
- [13] R. HARVEY & B. LAWSON: *A theory of characteristic currents associated with a singular connection*, Astérisque No. 213 (1993), 268 pp.
- [14] R. HARVEY & B. LAWSON: *Geometric residue theorems*, Amer. J. Math. **117** (1995), 829–873.
- [15] R. HARVEY & B. LAWSON: *Singularities and Chern-Weil theory. II. Geometric atomicity*, Duke Math. J. **119** (2003), 119–158.
- [16] M. MEO: *Résidus dans le cas non nécessairement intersection complète*, C. R. Acad. Sci. Paris Sér I Math. **333** (2001), 33–38.
- [17] M. MEO: *Courants résidus et formule de King*, Preprint (2003); to appear in Arkiv mat. .
- [18] M. PASSARE & A. TSIKH & A. YGER: *Residue currents of the Bochner-Martinelli type*, Publ. Mat. **44** (2000), 85–117.

DEPARTMENT OF MATHEMATICS, CHALMERS UNIVERSITY OF TECHNOLOGY  
 AND THE UNIVERSITY OF GÖTEBORG, S-412 96 GÖTEBORG, SWEDEN  
*E-mail address:* matsa@math.chalmers.se